Continuous Glucose Monitor (CGM)

| | | lication Information | |
|--|--|--|--|
| *Member ID: | " indicates | *Member Name: | |
| *DOB: | | *Weight: | |
| *Medication Name/Strength: | | ☐ Do Not Substitute. Authorizations will be processed for | |
| *Directions for use: | | the preferred Generic/B | rand equivalent unless specified |
| | Provider | Information | |
| | | required field | |
| *Requesting Provider Name: | | *NPI: | |
| *Address: | | | |
| *Contact Person: | | *Phone #: | |
| *Fax #: | | Email: | |
| | | g: laboratory results, chart not | • |
| provide | letter to Pharmacy PA at 855 | -828-4992 , to prevent process | ing delays. |
| Dexcom G6 and G7 *Preferred Product | Freestyle Libre 2 and Libre 3 | Freestyle Libre 14 days | Guardian Connect |
| 2 years of age and olde | er 4 years of age and older | 18 years of age and older | 14 to 75 years of age |
| ☐ Patient and/or cares | cognize and respond to the mess t the patient and/or caregiver hav | Gestational e diabetes treatment plan supervages, alarms, and alerts of the developed (or will receive) appropriation | evice. |
| (All the following criteria) Patient has been ad One of the following Patient's insulir Patient has hypewarning symptom Patient experies are not attributed A history of one mental and/or | herent to blood glucose testing. g applies: n regimen requires frequent adjuted by the programment of the pr | stment based on BGM or CGM te of neuroglycopenia before the app ant fall in blood glucose below no 2 hypoglycemia (glucose level of l cose level less than 54 mg/dl char ty assistance for treatment of hyp | pearance of autonomic rmal levels). ess than 54 mg/dl), which racterized by altered |
| Patient rPatient i | ng criteria must be met: equires insulin OR s not insulin-dependent and has nical rationale for CGM: | a history of problematic hypoglyo | cemia. |

Page 1 of 2 Last Updated 5/1/2023

UTAH MEDICAID PHARMACY PRIOR AUTHORIZATION REQUEST FORM

| | itional Criteria for Non-Diabetes Endocrine Disorders Causing Glycemic Variability, Off-Label Use: | | | | |
|------|--|--|--|--|--|
| | Requests for any off-label indications must be supported by at least one (1) major multi-site study or three (3) smaller studies published in JAMA, NEJM, Lancet or other peer review specialty medical journals within the most | | | | |
| | recent five (5) years. Supporting documentation must be included. Compendia use must be recommended by | | | | |
| | generally accepted compendia such as American Hospital Formulary Service Drug Information (AHFS), United | | | | |
| | States Pharmacopeia-Drug Information (USP-DI), and the DRUGDEX Information System. | | | | |
| | Diagnosis: Duration of treatment: Chart Note Page #: | | | | |
| Repl | acement Receiver (May be authorized when documentation confirms all the following): | | | | |
| | | | | | |
| | Patient is compliant with device and the device is required and continues to provide benefit to the patient's diabetic regimen. | | | | |
| | Replacement cannot be obtained through the supplier or manufacturer (warranty has expired). | | | | |
| Non | -Preferred Product: (Criteria above must also be met) | | | | |
| | Trial and failure of preferred Dexcom G6 or G7, or prescriber must demonstrate medical necessity for non- preferred product. | | | | |
| | Details: Chart Note Page #: | | | | |
| Re-a | uthorization Criteria (All the following criteria must be met) | | | | |
| | Updated documentation from the treating provider indicating the device is required and continues to provide | | | | |
| _ | benefit to the patient's diabetic regimen. | | | | |
| | Documentation of a face-to-face visit with the provider in the last 6 months. | | | | |
| Ini | itial Authorization: Up to six (6) months | | | | |
| Re | -authorization: Up to one (1) year | | | | |
| PR | OVIDER CERTIFICATION | | | | |
| ۱h | ereby certify this treatment is indicated, necessary and meets the guidelines for use. | | | | |
| | | | | | |

Date

Prescriber's Signature

Page 2 of 2 Last Updated 5/1/2023